

RATLIFF-RUSH FILTRATIONS ASSOCIATED WITH IDEALS AND MODULES OVER A NOETHERIAN RING

TONY J. PUTHENPURAKAL AND FAHED ZULFEQARR

ABSTRACT. Let R be a commutative Noetherian ring, M a finitely generated R -module and I a proper ideal of R . In this paper we introduce and analyze some properties of $r(I, M) = \bigcup_{k \geq 1} (I^{k+1}M : I^kM)$, the *Ratliff-Rush ideal associated with I and M* . When $M = R$ (or more generally when M is projective) then $r(I, M) = \tilde{I}$, the usual Ratliff-Rush ideal associated with I . If I is a regular ideal and $\text{ann } M = 0$ we show that $\{r(I^n, M)\}_{n \geq 0}$ is a stable I -filtration. If $M_{\mathfrak{p}}$ is free for all $\mathfrak{p} \in \text{Spec } R \setminus \text{m-Spec } R$, then under mild condition on R we show that for a regular ideal I , $\ell(r(I, M)/\tilde{I})$ is finite. Further $r(I, M) = \tilde{I}$ if $A^*(I) \cap \text{m-Spec } R = \emptyset$ (here $A^*(I)$ is the stable value of the sequence $\text{Ass}(R/I^n)$). Our generalization also helps to better understand the usual Ratliff-Rush filtration. When I is a regular $\mathfrak{m}$ -primary ideal our techniques yield an easily computable bound for k such that $\tilde{I}^n = (I^{n+k} : I^k)$ for all $n \geq 1$. For any ideal I we show that $\widetilde{I^n M} = I^n M + H_I^0(M)$ for all $n \gg 0$. This yields that $\tilde{\mathcal{R}}(I, M) = \bigoplus_{n \geq 0} \widetilde{I^n M}$ is Noetherian if and only if $\text{depth } M > 0$. Surprisingly if $\dim M = 1$ then $\tilde{G}_I(M) = \bigoplus_{n \geq 0} \widetilde{I^n M / I^{n+1} M}$ is always a Noetherian and a Cohen-Macaulay $G_I(R)$ -module. Application to Hilbert coefficients is also discussed.

INTRODUCTION

Let R be a commutative Noetherian ring and I an ideal of R . The Ratliff-Rush ideal $\tilde{I} = \bigcup_{k \geq 1} (I^{k+1} : I^k)$ is a useful notion. When R is local and I is $\mathfrak{m}$ -primary, the Ratliff-Rush filtration $\{\tilde{I}^n\}_{n \geq 1}$ has many applications in the theory of Hilbert functions, for instance see [18]. In this paper we generalize this notion.

Let M be a finitely generated R -module. Set

$$r(I, M) = \bigcup_{k \geq 1} (I^{k+1}M : I^kM).$$

We call $r(I, M)$ the *Ratliff-Rush ideal associated with I and M* . Notice that $r(I, M) = \tilde{I}$ if $M = R$. Our generalization also gives us a better understanding of the usual Ratliff-Rush filtration. For instance when R is a Cohen-Macaulay local ring and I is an $\mathfrak{m}$ -primary ideal it is useful find a upper bound on k such that $\tilde{I}^n = (I^{n+k} : I^k)$ for all $n \geq 1$, see [19]. Our techniques enables us to find an easily computable upper bound on k (see 8.12 and 8.7).

Date: February 2, 2008.

1991 *Mathematics Subject Classification.* Primary 13A15; Secondary 13D40, 13A30.

Key words and phrases. Ratliff-Rush Filtration, reductions, asymptotic associated primes, Hilbert functions.

The first author was partly supported by IIT Bombay seed grant 03ir053. The second author thanks CSIR for financial support.

We analyze many of its properties. Perhaps the first non-trivial property is that the function $I \mapsto r(I, M)$ is an involution on the set of ideals of R , i.e.,

$$r(r(I, M), M) = r(I, M).$$

This is done in Theorem 3.3.

Next we relate this notion to integral closure. In Theorem 4.4 we show that if I is a regular ideal then there exists a rank 1 module M such that $r(I, M) = \bar{I}$. A typical example of a rank one module is a regular ideal. In Proposition 4.5 we show that there exists a regular ideal J such that $r(I, J) = \bar{I}$. Furthermore we prove (in Proposition 4.5) that the set

$$\mathcal{C}(I) = \{J \mid J \text{ a regular ideal with } r(I, J) = \bar{I}\}.$$

has a unique maximal element.

Next we analyze the filtration $\mathcal{F}_M^I = \{r(I^n, M)\}_{n \geq 1}$. We first prove that this is a filtration of ideals and an I -filtration (see Theorem 2.1). Thus $\mathcal{R}(\mathcal{F}_M^I) = \bigoplus_{n \geq 0} r(I^n, M)$ is a $\mathcal{R}(I) = \bigoplus_{n \geq 0} I^n$ -algebra. Let $\widetilde{IM} = \bigcup_{k \geq 1} (I^{k+1}M :_M I^k)$, the Ratliff-Rush module of M associated with I . We show that $\widetilde{\mathcal{R}}(I, M) = \bigoplus_{n \geq 0} \widetilde{I^n M}$ is a graded $\mathcal{R}(\mathcal{F}_M^I)$ -module (see Proposition 2.4). In Theorem 5.3 we prove that if $\text{grade}(I, M) > 0$ and $\text{ann } M = 0$ then $\mathcal{F}_M^I$ is a stable I -filtration.

If M is a projective R -module then $r(I, M) = \widetilde{I}$ for all ideals I (see Proposition 1.6). As R is Noetherian, projective (finitely generated) modules are precisely locally free (finitely generated) modules. In some sense the next case is to consider R -modules M such that

$$(*) \quad M_{\mathfrak{p}} \text{ is free for all } \mathfrak{p} \in \text{Spec}(R) \setminus \text{m-Spec}(R)$$

(here $\text{m-Spec } R$ denotes the set of maximal ideals of R). For instance if $M \subset F$ where F is a free R -module and if $\ell(F/M)$ is finite then M satisfies (*). Here $\ell(-)$ denotes the length. On R we impose a mild condition $\text{Ass } R \cap \text{m-Spec } R = \emptyset$ (every domain that is not a field satisfies this condition). We show that if I is a regular ideal then $\ell(r(I, M)/\widetilde{I})$ is finite and the function $n \mapsto \ell(r(I^n, M)/\widetilde{I^n})$ is a polynomial function. We show that $r(I, M) = \widetilde{I}$ if $A^*(I) \cap \text{m-Spec } R = \emptyset$. Here $A^*(I)$ is the stable value of $\text{Ass}(R/I^n)$.

In the case when $\text{ann } M \neq 0$ or $\text{grade}(I, M) = 0$ we go modulo the ideal $(H_I^0(M) : M)$. In Proposition 7.3 we show that if $M \neq H_I^0(M)$ then

$$r(I^{n+1}, M) = I \cdot r(I^n, M) + (H_I^0(M) : M) \quad \text{for all } n \gg 0.$$

An easy consequence of our techniques (see Proposition 7.5) is the following result in the case of the usual Ratliff-Rush filtration of a module M with respect to I

$$\widetilde{I^n M} = I^n M + H_I^0(M) \quad \text{for all } n \gg 0.$$

In the final section we show that if $\dim M = 1$, $\widetilde{G}_I(M) = \bigoplus_{n \geq 0} \widetilde{I^n M} / \widetilde{I^{n+1} M}$ is a Noetherian and a Cohen-Macaulay $G_I(R)$ -module. This is surprising since if $\text{depth } M = 0$ (and $\dim M = 1$) then $\widetilde{\mathcal{R}}(I, M) = \bigoplus_{n \geq 0} \widetilde{I^n M}$ is not a Noetherian $\mathcal{R}(I)$ -module. We also give an application of our result to Hilbert coefficients.

Here is an overview of the contents of the paper. In section one we study few basic properties of the ideal $r(I, M)$. In section two we study the filtration $\{r(I^n, M)\}_{n \geq 1}$ and explore the relation between $\{r(I^n, M)\}_{n \geq 1}$ and $\{\widetilde{I^n M}\}_{n \geq 1}$. In section three

we prove that the operation $I \mapsto r(I, M)$ is an involution. In section four we relate it to integral closure. In section five we prove that it is a stable I -filtration when I is regular and $\text{ann } M = 0$. In section six we study the case when $M_{\mathfrak{p}}$ is free for all $\mathfrak{p} \in \text{Spec } R \setminus \text{m-Spec } R$. In section seven we study the general case when $\text{ann } M \neq 0$ or $\text{grade}(I, M) = 0$. For the next sections we assume that $(R, \mathfrak{m})$ is local. In section eight we study its relation with superficial elements. This is then used to give a bound on k such that $\widetilde{I}^n = (I^{n+k} : I^k)$ for all $n \geq 1$. In section nine we study ideals having a principal reduction and use it to compute $r(I, M)$ in some examples. In final section we show that if $\dim M = 1$ then $\widetilde{G}_I(M) = \bigoplus_{n \geq 0} \widetilde{I}^n M / \widetilde{I}^{n+1} M$ is always a Cohen-Macaulay $G_I(R)$ -module. Finally we give an application of one of our results to Hilbert coefficients of a 1-dimensional module.

1. PRELIMINARIES

In this paper unless otherwise stated all rings considered are commutative Noetherian and all modules are assumed to be finitely generated.

Let R be a ring and I an ideal of R . Let M be an R -module. Consider the following ascending chain of ideals in R

$$I \subseteq (IM : M) \subseteq (I^2M : IM) \subseteq \dots \subseteq (I^{k+1}M : I^kM) \subseteq \dots$$

Since R is Noetherian, this chain stabilizes. We denote its stable value by $r(I, M)$. We call $r(I, M)$ the *Ratliff-Rush ideal associated with I and M* .

In this section we prove some basic properties of ideal $r(I, M)$, in particular we show that

$$r(I, M \oplus N) = r(I, M) \cap r(I, N) \quad \text{and} \quad r(I, M \otimes_R N) \supseteq r(I, M) + r(I, N).$$

We also investigate the case when $r(I, M) = R$.

Remarks 1.1.

- (a) $r(I, M) = (I^{k+1}M : I^kM)$ for all $k \gg 0$.
- (b) When $M = R$,

$$r(I, R) = \bigcup_{k \geq 1} (I^{k+1} : I^k) = \widetilde{I}, \text{ the Ratliff-Rush closure of } I. \quad (\text{see [11]})$$

- (c) For $n \geq 1$, we have

$$r(I^n, M) = \bigcup_{k \geq 1} (I^{n+k}M : I^kM).$$

- (d) One can easily check that

$$I \subseteq \widetilde{I} \subseteq r(I, M).$$

- (e) Let $R = \bigoplus_{n \geq 0} R_n$ be a graded ring and I , a homogeneous ideal of R . Let M be a graded R -module. Then $r(I, M)$ is a homogeneous ideal.

We give an example which shows that there exists a module M such that

$$I \subsetneq \widetilde{I} \subsetneq r(I, M).$$

Example 1.2. Let $R = k[t^4, t^{11}, t^{17}, t^{18}]$, $\mathfrak{m} = \langle t^4, t^{11}, t^{17}, t^{18} \rangle R$, $I = \langle t^4, t^{11} \rangle R$ and $M = \langle t^4, t^{11}, t^{17} \rangle R$. One can prove (by induction on n) that

$$I^n = \langle t^{4n+7i} : i = 0, 1, 2, \dots, n \rangle R \quad \text{for all } n \geq 1.$$

Note that $t^{18} \cdot I^2 \subseteq I^3$ so $t^{18} \in \tilde{I}$. One can verify that $t^{17} \notin (I^{n+1} : I^n)$ for all $n \geq 1$. Thus we get $\tilde{I} = \langle t^4, t^{11}, t^{18} \rangle R$. Notice that $t^{17} \in (IM : M)$. Now since $\tilde{I} \subseteq r(I, M)$, we get $r(I, M) = \langle t^4, t^{11}, t^{17}, t^{18} \rangle R$. Therefore $I \subsetneq \tilde{I} \subsetneq r(I, M)$ and $r(I, M) = \mathfrak{m}$.

We next give an example from [17, 1.4] of a Ratliff-Rush closed ideal I of ring R and a module M such that $I \neq r(I, M)$.

Example 1.3. Let $R = k[x, y]$ be a polynomial ring in variables x and y . Let

$$\begin{aligned} I &= \langle y^{22}, x^4 y^{18}, x^7 y^{15}, x^8 y^{14}, x^{11} y^{11}, x^{14} y^8, x^{15} y^7, x^{18} y^4, x^{22} \rangle R, \\ \text{and } M &= \langle x^2, y^3 \rangle R. \end{aligned}$$

By [17, 1.4], I is Ratliff-Rush closed, that is $I = \tilde{I}$ (also see Example 8.13). Using *Singular* [7] one can verify that

$$(I^4 M : I^3 M) = I + \langle x^2 y^{21}, x^6 y^{17}, x^{13} y^{10}, x^{20} y^3 \rangle R.$$

So, $I \subsetneq (I^4 M : I^3 M) \subseteq r(I, M)$. Hence $I = \tilde{I} \subsetneq r(I, M)$.

We collect some properties of the ideal $r(I, M)$ in the following proposition.

Proposition 1.4. *Let M and N be any two R -modules. Then*

- (a) *If $f : M \rightarrow N$ is a surjective R -homomorphism then $r(I, M) \subseteq r(I, N)$.*
- (b) *$r(I, M \oplus N) = r(I, M) \cap r(I, N)$.*
- (c) *$r(I, M \otimes_R N) \supseteq r(I, M) + r(I, N)$.*
- (d) *If T is a Noetherian ring which is a flat R -algebra then*

$$r(I, M) \otimes_R T = r_T(IT, M \otimes_R T).$$

- (e) *If S is a multiplicative closed subset of R then*

$$r(I, M) \otimes_R R_S = r_{R_S}(IR_S, M_S).$$

- (f) *For a regular ideal I , $r(\tilde{I}, M) = r(I, M)$.*
- (g) *For each $n \geq 1$, we have $r(I^n, M) = r(I^n, I^s M)$ for all $s \geq 1$.*

Proof. (a) Let $x \in r(I, M)$. Then, for some $k \geq 0$, we have $x \in (I^{k+1} M : I^k M)$. Thus

$$\begin{aligned} x I^k M &\subseteq I^{k+1} M, \\ \text{so } x I^k f(M) &\subseteq I^{k+1} f(M). \end{aligned}$$

Since f is surjective, we get $x \in (I^{k+1} N : I^k N) \subseteq r(I, N)$.

(b) Let $x \in r(I, M \oplus N)$. Then $x I^k (M \oplus N) \subseteq I^{k+1} (M \oplus N)$ for some $k \geq 0$. Therefore

$$x I^k M \subseteq I^{k+1} M \quad \text{and} \quad x I^k N \subseteq I^{k+1} N.$$

$$\text{So } x \in r(I, M) \cap r(I, N).$$

The reverse inclusion is obvious.

(c) Let $x \in r(I, M)$. We have $x I^k M \subseteq I^{k+1} M$ for some $k \geq 0$. Let $\sum_i \alpha_i (m_i \otimes n_i) \in I^k (M \otimes N)$, where $\alpha_i \in I^k$. We have

$$x \left(\sum_i \alpha_i (m_i \otimes n_i) \right) = \sum_i (x \alpha_i m_i) \otimes n_i$$

Since $x\alpha_i m_i \in I^{k+1}M$, $x\alpha_i m_i = \sum_j \beta_j^{(i)} u_j^{(i)}$, where $\beta_j^{(i)} \in I^{k+1}$ and $u_j^{(i)} \in M$. Thus

$$\begin{aligned} x \left(\sum_i \alpha_i (m_i \otimes n_i) \right) &= \sum_i \left(\sum_j \beta_j^{(i)} u_j^{(i)} \right) \otimes n_i \\ &= \sum_{i,j} (\beta_j^{(i)} u_j^{(i)}) \otimes n_i \\ &= \sum_{i,j} \beta_j^{(i)} (u_j^{(i)} \otimes n_i) \in I^{k+1}(M \otimes N). \end{aligned}$$

In the same way one can show that $r(I, N) \subseteq r(I, M \otimes_R N)$.

(d) Using [14, 18.1], we have

$$(I^{n+1}M : I^n M) \otimes_R T = (I^{n+1}M \otimes_R T :_T I^n M \otimes_R T) \quad \text{for all } n \geq 1.$$

Therefore we get

$$\begin{aligned} r(I, M) \otimes_R T &= (I^{k+1}M : I^k M) \otimes_R T \quad \text{for all } k \gg 0, \\ &= (I^{k+1}M \otimes_R T :_T I^k M \otimes_R T) \quad \text{for all } k \gg 0, \\ &= r_T(IT, M \otimes_R T). \end{aligned}$$

(e) This follows from part (d), since R_S is a flat R -algebra.

(f) For any regular ideal I of R , we have $I^k = \tilde{I}^k$ for all $k \gg 0$ (see [11, 2.1]). Therefore

$$\begin{aligned} r(\tilde{I}, M) &= (\tilde{I}^{k+1}M : \tilde{I}^k M) \quad \text{for all } k \gg 0, \\ &= (I^{k+1}M : I^k M) \quad \text{for all } k \gg 0, \\ &= r(I, M). \end{aligned}$$

(g) Notice that

$$r(I^n, I^s M) = \bigcup_{k \geq 1} (I^{n+k} I^s M : I^k I^s M) = r(I^n, M).$$

□

Remark 1.5. From 1.4(b) and 1.1(d) it follows that if a module M has a free summand then $r(I, M) = \tilde{I}$.

Corollary 1.6. *Let M be a projective R -module and let I be an ideal of R . Then $r(I, M) = \tilde{I}$.*

Proof. Set $J = r(I, M)$. Since M is projective, $M_{\mathfrak{p}}$ is free for all $\mathfrak{p} \in \text{Spec } R$, by [5, 4.11 (b)]. Note that $\tilde{I} \subseteq J$. Now if $\mathfrak{p} \in \text{Spec } R$ then

$$\begin{aligned} J_{\mathfrak{p}} &= r_{R_{\mathfrak{p}}}(I_{\mathfrak{p}}, M_{\mathfrak{p}}), \quad (\text{by 1.4e}) \\ \text{Also } (\tilde{I})_{\mathfrak{p}} &= \tilde{I}_{\mathfrak{p}}. \end{aligned}$$

Thus $J/\tilde{I}$ is locally zero and hence zero. □

The next proposition enables us to determine when is $r(I, M) = R$. Let

$$H_I^0(M) = \{m \in M : I^n m = 0 \text{ for some } n \geq 1\}$$

be the I -torsion submodule of M .

Proposition 1.7. *Let I be a proper ideal of ring R . The following conditions are equivalent*

- (a) $r(I, M) = R$.
- (b) *there exists $n \geq 0$ such that $I^n M = I^{n+1} M$.*

If $H_I^0(M) = M$ then $r(I, M) = R$. Furthermore if R is local then the converse is also true.

Proof. (a) $\Rightarrow$ (b): If $r(I, M) = R$ then $1 \in (I^{n+1} M : I^n M)$ for some n . Therefore we get $I^{n+1} M = I^n M$.

(b) $\Rightarrow$ (a): If $I^n M = I^{n+1} M$ for some $n \geq 0$ then $1 \in (I^{n+1} M : I^n M) \subseteq r(I, M)$. So $r(I, M) = R$.

Suppose $H_I^0(M) = M$. Since M is a finitely generated R -module, there exists an integer $n \geq 1$ such that $I^n M = 0$. Hence $r(I, M) = R$.

Let R be local and $I \subseteq \mathfrak{m}$. If $r(I, M) = R$ then by (b), $I^n M = I^{n+1} M$. So by Nakayama's lemma, $I^n M = 0$. Thus $H_I^0(M) = M$. $\square$

The following example shows that in the non-local case, it is possible that $r(I, M) = R$ but $H_I^0(M) \neq M$.

Example 1.8. Let R be a ring, having a nontrivial idempotent element e (i.e. $e^2 = e$ and $e \neq 0, 1$). Let $I = \langle e \rangle R$ and $M = I$. Notice that $I^n M = I^{n+1} M$ for all $n \geq 1$. Therefore, by Proposition 1.7, we have $r(I, M) = R$. Note that e cannot be killed by any power of I as $I^n \cdot e = I^{n+1} = I \neq 0$. So $e \notin H_I^0(M)$. But $e \in M$. Hence $M \neq H_I^0(M)$.

2. THE FILTRATION $\mathcal{F}_M^I = \{r(I^n, M)\}_{n \geq 0}$

This section deals with $\mathcal{F}_M^I = \{r(I^n, M)\}_{n \geq 0}$. We first show that it is a filtration of ideals and also an I -filtration. We explore its relation with the Ratiff-Rush filtration of M with respect to I . We also prove that if $\text{grade}(I, M) > 0$ then $r(I, M) = (\tilde{I}M : M)$.

For the definition of filtration of ideals, see [3, 4.4]. The following theorem shows that the collection $\mathcal{F}_M^I = \{r(I^n, M)\}_{n \geq 0}$ of ideals is an I -filtration.

Theorem 2.1. *For any R -module M , the sequence $\mathcal{F}_M^I$ is a filtration of ideals. It is also an I -filtration.*

Proof. It is easy to show that

$$r(I^n, M) \supseteq r(I^{n+1}, M) \quad \text{for all } n \geq 0.$$

Next we prove the following

- (a) $r(I^n, M) \cdot r(I^m, M) \subseteq r(I^{n+m}, M)$ for all $n, m \geq 1$.
 - (b) $I \cdot r(I^n, M) \subseteq r(I^{n+1}, M)$ for all $n \geq 1$.
- (a): Let $x \in r(I^n, M)$ and $y \in r(I^m, M)$. We have

$$\begin{aligned} xI^k M &\subseteq I^{n+k} M \quad \text{for all } k \gg 0, \\ \text{and } yI^k M &\subseteq I^{m+k} M \quad \text{for all } k \gg 0. \end{aligned}$$

Therefore, for all $k \gg 0$, we have

$$xyI^k M \subseteq xI^{m+k} M \subseteq I^{n+m+k} M.$$

$$\text{Thus, } xy \in r(I^{n+m}, M).$$

(b): This follows from (a) since $I \subseteq r(I, M)$. Thus $\mathcal{F}_M^I$ is an I -filtration. $\square$

Let us recall the definition (see [9], also see [15]) of the Ratliff-Rush submodule of M with respect to I .

Definition 2.2. Consider the following chain of submodules of M :

$$IM \subseteq (I^2M :_M I) \subseteq (I^3M :_M I^2) \subseteq \dots \subseteq (I^{n+1}M :_M I^n) \subseteq \dots$$

Since M is Noetherian, this chain of submodules stabilizes. We denote its stable value by $\widetilde{IM}$. We call $\widetilde{IM}$ to be the *Ratliff-Rush submodule of M associated with I* . The filtration $\{\widetilde{I^n M}\}_{n \geq 1}$ is called the *Ratliff-Rush filtration of M with respect to I* .

Notation: To facilitate further calculations, set

$$\begin{aligned} \mathcal{F}_M^I &= \{r(I^n, M)\}_{n \geq 0}, & \mathcal{R}(\mathcal{F}_M^I) &= \bigoplus_{n \geq 0} r(I^n, M), \\ \mathcal{R}(I) &= \bigoplus_{n \geq 0} I^n, & G_I(R) &= \bigoplus_{n \geq 0} I^n / I^{n+1}, & G_I(R)_+ &= \bigoplus_{n \geq 1} I^n / I^{n+1}, \\ G_I(M) &= \bigoplus_{n \geq 0} I^n M / I^{n+1} M & \text{and} & & \widetilde{\mathcal{R}}(I, M) &= \bigoplus_{n \geq 0} \widetilde{I^n M}. \end{aligned}$$

Remark 2.3. By Theorem 2.1, $\mathcal{F}_M^I$ is a filtration of ideals in R , so $\mathcal{R}(\mathcal{F}_M^I)$ is a ring. Clearly $\mathcal{R}(I) \subseteq \mathcal{R}(\mathcal{F}_M^I)$ is a subring. Since $\mathcal{F}_M^I$ is an I -filtration then $\mathcal{R}(\mathcal{F}_M^I)$ is an $\mathcal{R}(I)$ -module.

We study the relation between the $\mathcal{R}(I)$ -algebra $\mathcal{R}(\mathcal{F}_M^I)$ and the $\mathcal{R}(I)$ -module $\widetilde{\mathcal{R}}(M)$. We first show that

Proposition 2.4. $\widetilde{\mathcal{R}}(I, M)$ is a graded $\mathcal{R}(\mathcal{F}_M^I)$ -module.

Proof. Set $I_n = r(I^n, M)$. It is enough to check that

$$(*) \quad I_n \cdot \widetilde{I^m M} \subseteq \widetilde{I^{n+m} M} \quad \text{for all } n, m.$$

Take any $x \in I_n$ and $z \in \widetilde{I^m M}$. Then, by [15, 2.2(iii)] and definition of I_n , we have

$$I^k z \subseteq I^{k+m} M \quad \text{and} \quad x I^k M \subseteq I^{k+n} M \quad \text{for all } k \gg 0.$$

Therefore, for all $k \gg 0$

$$\begin{aligned} x I^k z &\subseteq x \cdot I^{m+k} M \subseteq I^{n+m+k} M. \\ \text{So } xz &\in \widetilde{I^{n+m} M}. \end{aligned}$$

□

Corollary 2.5. Set $J = r(I, M)$. Then

$$I^n M \subseteq J^n M \subseteq \widetilde{I^n M} \quad \text{for all } n \geq 1.$$

Proof. The first inclusion is clear. The second inclusion follows from the fact $J^n \subseteq r(I^n, M)$ and Proposition 2.4. □

Corollary 2.6. Set $J = r(I, M)$. Assume that $\text{grade}(I, M) > 0$. Then

$$J^n M = I^n M \quad \text{for all } n \gg 0.$$

Furthermore if $\text{grade}(G_I(R)_+, G_I(M)) > 0$ then

$$J^n M = I^n M \quad \text{for all } n \geq 1,$$

Proof. First part follows from [15, 2.2]. For second part, note that $\text{grade}(G_I(R)_+, G_I(M)) > 0$ implies (see [9, Fact 9]) that $\widetilde{I^n M} = I^n M$ for all $n \geq 1$. $\square$

Proposition 2.7. *If $\text{grade}(I, M) > 0$ then*

$$r(I, M) = (\widetilde{IM} : M).$$

Proof. From Proposition 2.4, it follows that $r(I, M) \subseteq (\widetilde{IM} : M)$. For reverse inclusion, let $x \in (\widetilde{IM} : M)$. Then

$$\begin{aligned} xM &\subseteq \widetilde{IM}, \\ \text{so } xI^k M &\subseteq I^k \widetilde{IM} = I^{k+1} M \text{ for all } k \gg 0. \end{aligned}$$

Thus $x \in r(I, M)$. Hence $r(I, M) = (\widetilde{IM} : M)$. $\square$

3. INVOLUTION PROPERTIES

In this section we prove that the function $I \mapsto r(I, M)$ is an involution on the set of ideals of R (see Theorem 3.3). We first prove the result in the case when $\text{grade}(I, M) > 0$. We also show that if $\text{grade}(I, M) > 0$ then $r(I, M)$ is a Ratliff-Rush closed ideal.

Proposition 3.1. *Let $\text{grade}(I, M) > 0$. Set $J = r(I, M)$. Then*

- (a) $r(J, M) = J$.
- (b) $\widetilde{J} = J$.

Proof. (a) By Corollary 2.6, there exists an integer k_0 such that

$$(i) \quad J^k M = I^k M \text{ for } k \geq k_0.$$

Also there exists k_0' such that

$$r(J, M) = (J^{k+1} M : J^k M) \text{ for all } k \geq k_0'.$$

From (i), it follows that

$$\begin{aligned} r(J, M) &= (I^{k+1} M : I^k M) \text{ for all } k \geq \max\{k_0, k_0'\} \\ &\subseteq r(I, M) = J. \end{aligned}$$

Since $J \subseteq r(J, M)$ always we get $r(J, M) = J$.

(b) Let $x \in \widetilde{J}$. Then

$$\begin{aligned} xJ^k &\subseteq J^{k+1} \text{ for all } k \gg 0, \\ xJ^k M &\subseteq J^{k+1} M \text{ for all } k \gg 0. \end{aligned}$$

By Corollary 2.6, we get

$$xI^k M \subseteq I^{k+1} M \text{ for all } k \gg 0.$$

Therefore $x \in J$. Hence $\widetilde{J} = J$. $\square$

To tackle the case when $\text{grade}(I, M) = 0$ we first prove

Lemma 3.2. *For any ideal I , the following hold*

- (a) $r(I, M) = r(I, M/H_I^0(M))$.
- (b) $r(I^n, M) = r(I^n, M/H_I^0(M))$ for all $n \geq 1$.

Proof. (a) If $M = H_I^0(M)$ then our assertion follows from Proposition 1.7. Suppose $M \neq H_I^0(M)$. Since the natural map $M \longrightarrow M/H_I^0(M)$ is surjective, by Proposition 1.4(a), we have

$$r(I, M) \subseteq r(I, M/H_I^0(M)).$$

Let $x \in r(I, M/H_I^0(M))$. Therefore

$$xI^k M + H_I^0(M) \subseteq I^{k+1}M + H_I^0(M) \quad \text{for some } k \geq 0.$$

Since M is a finitely generated R -module, there exists an integer $r \in \mathbb{N}$ such that $I^r H_I^0(M) = 0$. Therefore

$$xI^{r+k}M \subseteq I^{r+k+1}M.$$

So $x \in (I^{r+k+1}M : I^{r+k}M) \subseteq r(I, M)$.

(b) This follows from (a), since $H_{I^n}^0(M) = H_I^0(M)$ for all $n \geq 1$. $\square$

We now prove the involution property in general.

Theorem 3.3. *For any ideal I of ring R , we have*

$$r(r(I, M), M) = r(I, M).$$

Proof. Set $J = r(I, M)$ and $N = M/H_I^0(M)$. If $M = H_I^0(M)$, then $J = R$ by Proposition 1.7. Also clearly $r(R, M) = R$. Now suppose $M \neq H_I^0(M)$. By using Lemma 3.2 for the ideals I and J , we have

$$(i) \quad J = r(I, M) = r(I, N).$$

$$(ii) \quad r(J, M) = r(J, M/H_J^0(M)).$$

Since $I \subseteq J$ we get $H_J^0(M) \subseteq H_I^0(M)$. Therefore the epimorphism

$$M/H_J^0(M) \longrightarrow M/H_I^0(M) = N$$

induces, by Proposition 1.4(a),

$$(iii) \quad r(J, M/H_J^0(M)) \subseteq r(J, N).$$

Notice that $\text{grade}(I, N) > 0$. Therefore, from Proposition 3.1, we get

$$(iv) \quad r(J, N) = r(r(I, N), N) = r(I, N).$$

Using (ii), (iii) and (iv), we get

$$r(J, M) = r(J, M/H_J^0(M)) \subseteq r(J, N) = J.$$

But $r(J, M) \supseteq J$ always. Therefore $r(J, M) = J$. $\square$

4. RELATION WITH INTEGRAL CLOSURE

In this section we show that $r(I, M) \subseteq \bar{I}$, the integral closure of I when $\text{ann } M = 0$ and $\text{grade}(I, R) > 0$. In particular when M has a positive rank and I is a regular ideal. We prove that if I is a regular ideal then there exists an R -module M of rank 1 such that $r(I, M) = \bar{I}$. Finally we show that if I is a regular ideal then the set

$$\mathcal{C}(I) := \{ J : J \text{ is regular ideal and } r(I, J) = \bar{I} \}$$

is non-empty and has a unique maximal element.

Remark 4.1. For any $x \in r(I, M)$, there exists $k \geq 0$ such that $xI^k M \subseteq I^{k+1}M$. By determinant trick, there exists $f(t) \in R[t]$ such that

$$f(t) = t^n + a_1 t^{n-1} + \dots + a_{n-1} t + a_n, \text{ where } a_i \in I^i,$$

and $f(x) \in \text{ann}(I^k M)$.

The following proposition gives a relation between $r(I, M)$ and $\bar{I}$.

Proposition 4.2. *Let $\text{ann } M = 0$. If either of following two conditions holds*

- (a) $\text{grade}(I, R) > 0$.
- (b) $\text{grade}(I, M) > 0$.

Then $r(I, M) \subseteq \bar{I}$. In particular if I is a regular ideal and M has a positive rank then $r(I, M) \subseteq \bar{I}$.

Proof. (a) By the remark above we get that $f(x)I^k M = 0$ for some $k \geq 0$. But $\text{ann } M = 0$. So $f(x) \cdot I^k = 0$. Since $\text{grade}(I, R) > 0$, we get $f(x) = 0$. Hence $x \in \bar{I}$. (b) Note that $\text{grade}(I, M) > 0$ yields $\text{ann}(I^k M) = \text{ann } M$. Now from hypothesis $\text{ann } M = 0$, it follows that $f(x) = 0$. Hence $x \in \bar{I}$.

Finally note that if M has a positive rank, say r , then $\text{ann } M = 0$ (since it contains R^r as a submodule). $\square$

Remark 4.3. Notice that $\text{ann } M = 0$ together with $\text{grade}(I, M) > 0$ implies $\text{grade}(I, R) > 0$. Thus (b) follows from (a).

The next proposition ensures the existence of an R -module M for a regular ideal I of R such that $r(I, M) = \bar{I}$.

Theorem 4.4. *Let I be a regular ideal of R . Let J be an ideal such that $I \subseteq J \subseteq \bar{I}$. Then there exists R -module M of rank 1 such that*

$$I \subseteq J \subseteq r(I, M) \subseteq \bar{I}.$$

In particular, there exists an R -module M of rank 1 such that $r(I, M) = \bar{I}$.

Proof. Let $z \in J \setminus I$ be any element. We have $z^n + \sum_{i=1}^n a_i z^{n-i} = 0$, with $a_i \in I^i$. Thus

$$(*) \quad z^n = - \sum_{i=1}^n a_i z^{n-i}.$$

Set $N = \langle z, I \rangle^{n-1} R$. We claim that $zN \subseteq IN$. Let $x \in N$. Then

$$x = rz^{n-1} + \sum_{i=1}^{n-1} b_i z^{n-1-i}, \text{ where } r \in R \text{ and } b_i \in I^i.$$

$$\text{Therefore } zx = rz^n + \sum_{i=1}^{n-1} b_i z^{n-i}.$$

By using (*), we get $zx = \sum_{i=1}^{n-1} (b_i - ra_i) z^{n-i} - ra_n \in IN$. Thus $zN \subseteq IN$ and hence $z \in r(I, N)$.

Let us assume that $J = \langle z_1, z_2, \dots, z_s \rangle R$ and set $N_i = \langle z_i, I \rangle^{n_i-1} R$, where n_i is the degree of an integral equation satisfied by z_i . Set $M = N_1 \otimes_R N_2 \otimes_R \dots \otimes_R N_s$. Notice that $\text{rank } M = 1$. By Proposition 1.4, we get $J \subseteq r(I, M)$. But, by Proposition 4.2(a), we always have $\bar{I} \supseteq r(I, M)$. $\square$

Proposition 4.5. *For any regular ideal I , there exists a regular ideal J such that*

$$r(I, J) = \bar{I}.$$

Proof. By Theorem 4.4, there exists an R -module M of rank 1 such that $r(I, M) = \bar{I}$. Let $T(M)$ denote the torsion submodule of M and set $N = M/T(M)$. Note that the surjective map $M \rightarrow N$ induces $r(I, N) \supseteq r(I, M) = \bar{I}$. But as $\text{rank } N = \text{rank } M = 1$, we have $r(I, N) \subseteq \bar{I}$ (by 4.2(a)). So $r(I, N) = \bar{I}$. Clearly N is torsion-free so that $N \simeq J$ for some ideal J of R . Since J is of rank 1, J is a regular ideal. Notice that $r(I, J) = r(I, N) = \bar{I}$. $\square$

Let $\mathcal{C}(I) := \{ J : J \text{ is a regular ideal and } r(I, J) = \bar{I} \}$. By Proposition 4.5, $\mathcal{C}(I) \neq \emptyset$. Since R is Noetherian, $\mathcal{C}(I)$ has a maximal element. We show that

Theorem 4.6. *$\mathcal{C}(I)$ has a unique maximal element.*

Proof. Suppose $Q \in \mathcal{C}(I)$ is a maximal element and $J \in \mathcal{C}(I)$. By Proposition 1.4, the following epimorphism $J \oplus Q \rightarrow J + Q \rightarrow 0$ induces

$$r(I, J \oplus Q) \subseteq r(I, J + Q).$$

$$\text{But } r(I, J \oplus Q) = r(I, J) \cap r(I, Q) = \bar{I}, \quad \text{by Proposition 1.4(b).}$$

$$\text{So } \bar{I} \subseteq r(I, J + Q).$$

But $r(I, J + Q) \subseteq \bar{I}$, since $J + Q$ has rank 1 as an R -module so $r(I, J + Q) = \bar{I}$. As Q is maximal this gives $Q = J + Q$. So $J \subseteq Q$. Hence $\mathcal{C}(I)$ has a unique maximal element. $\square$

5. STABLE FILTRATIONS

In this section we discuss the conditions under which our filtration $\mathcal{F}_M^I = \{r(I^n, M)\}_{n \geq 0}$ is a stable I -filtration. This is equivalent to saying that the Rees algebra $\mathcal{R}(\mathcal{F}_M^I)$ is a finitely generated $\mathcal{R}(I)$ -module. Our main result (Theorem 5.9) is that if $\text{grade}(I, R) > 0$ and $\text{ann } M = 0$ then $\mathcal{F}_M^I$ is a stable I -filtration. In local case we prove that $\text{ann } M = 0$ is a necessary condition for $\mathcal{F}_M^I$ to be a stable I -filtration.

5.1. Recall a filtration of ideals $R = I_0 \supseteq I_1 \supseteq \dots \supseteq I_n \supseteq I_{n+1} \supseteq \dots$ is said to be a *stable I -filtration* if $II_n \subseteq I_{n+1}$ for all $n \geq 0$ and $II_n = I_{n+1}$ for $n \gg 0$.

The lemma below is crucial to prove our main result.

Lemma 5.2. *Let S be a ring and $R \subseteq S$, a subring of S , such that R is Noetherian. Assume that there is a faithful S -module E (i.e., $\text{ann}_S(E) = 0$) such that E is a finitely generated R -module. Then S is finitely generated as a R -module (and so Noetherian).*

Proof. Note that any S -linear map $f : M \rightarrow N$ between S -modules M and N , is also R -linear. Consider the inclusion map

$$i : \text{Hom}_S(E, E) \rightarrow \text{Hom}_R(E, E) \quad \text{such that } f \mapsto f.$$

Notice that i is R -linear. For $s \in S$, let $\mu_s : E \rightarrow E$ be the multiplication map i.e., $\mu_s(t) = st$ for all $t \in E$. Define

$$\begin{aligned} \phi : S &\rightarrow \text{Hom}_S(E, E) \\ s &\mapsto \mu_s. \end{aligned}$$

Clearly ϕ is S -linear and so R -linear. Notice that $\ker \phi = 0$, since E is a faithful S -module. Consider the following composition

$$S \xrightarrow{\phi} \text{Hom}_S(E, E) \xrightarrow{i} \text{Hom}_R(E, E).$$

Clearly $i \circ \phi$ is an injective R -linear map. Therefore as R -modules

$$S \cong \text{ to a } R\text{-submodule of } \text{Hom}_R(E, E).$$

As R is Noetherian and E is a finitely generated R -module, we get S is a finitely generated R -module. $\square$

The next theorem shows that the filtration $\mathcal{F}_M^I$ is a stable I -filtration under fairly mild assumptions.

Theorem 5.3. *Let $\text{grade}(I, M) > 0$ and $\text{ann } M = 0$. Then $\mathcal{F}_M^I$ is a stable I -filtration.*

Proof. For convenience, set $\mathcal{S} = \mathcal{R}(\mathcal{F}_M^I)$, $\mathcal{R} = \mathcal{R}(I)$ and $E = \bigoplus_{n \geq 0} \widetilde{I^n M} t^n$. By Proposition 2.4, E is an $\mathcal{S}$ -module. Since $\text{grade}(I, M) > 0$, $\widetilde{I^n M} = I^n M$ for all $n \gg 0$ (see [15, 3.3]). So E is a finitely generated $\mathcal{R}$ -module.

We prove that $\text{ann}_{\mathcal{S}}(E) = 0$. Notice that $\text{ann}_{\mathcal{S}}(E)$ is a homogeneous ideal of $\mathcal{S}$. Let $xt^n \in \text{ann}_{\mathcal{S}}(E)$ be a homogeneous element. As $xt^n \cdot E = 0$ we get $x \cdot M = 0$. Thus $x \in \text{ann } M = 0$. Therefore $\text{ann}_{\mathcal{S}}(E) = 0$. Using Lemma 5.2, we conclude that S is a finitely generated $\mathcal{R}$ -module. So $\mathcal{F}_M^I$ is a stable I -filtration. $\square$

The following example shows that the hypothesis in Theorem 5.3 is not necessary for $\mathcal{R}(\mathcal{F}_M^I)$ to be Noetherian.

Example 5.4. Let I be a nilpotent ideal of R (i.e. $I^r = 0$ for some $r \geq 1$). Then $r(I^n, M) = R$ for all $n \geq 0$ and for any R -module M . Thus $\mathcal{R}(\mathcal{F}_M^I) \cong R[t]$, which is a Noetherian ring.

Remark 5.5. Let I be any ideal of R . Let $x \in \text{ann } M$. Then $x \cdot M = 0$, so $xI^k M = 0 \subseteq I^{n+k} M$ for all $n, k \geq 1$. Therefore $x \in r(I^n, M)$ for all $n \geq 1$. Hence

$$\text{ann } M \subseteq \bigcap_{n \geq 1} r(I^n, M).$$

Proposition 5.6. *If $\mathcal{F}_M^I$ is a stable I -filtration then*

$$\text{ann } M \subseteq \bigcap_{n \geq 1} I^n.$$

Proof. For convenience, set $I_n = r(I^n, M)$. Since $\mathcal{F}_M^I$ is a stable I -filtration, there exists an integer n_0 such that

$$(i) \quad I_{n_0+k} = I^k I_{n_0} \quad \text{for all } k \geq 1.$$

By above Remark 5.5, one has $\text{ann } M \subseteq I_{n_0+k}$ for all $k \geq 1$. Therefore, by (i), we have

$$\text{ann } M \subseteq I_{n_0+k} \subseteq I^k \quad \text{for all } k \geq 1.$$

Hence result follows. $\square$

Remark 5.7. The above Proposition 5.6 proves that if the I -adic filtration is *separated*, i.e. $\bigcap_{n \geq 1} I^n = 0$ then

$$\mathcal{F}_M^I \text{ is a stable } I\text{-filtration} \implies \text{ann } M = 0.$$

An easy consequence of Proposition 5.6 is following

Corollary 5.8. *Let $(R, \mathfrak{m})$ be a local ring and $M \neq H_I^0(M)$. Then*

$$\mathcal{F}_M^I \text{ is a stable } I\text{-filtration} \implies \text{ann } M = 0.$$

Proof. For a local ring $(R, \mathfrak{m})$, the I -adic filtration is separated (by Krull's intersection theorem). Hence $\text{ann } M = 0$. $\square$

In the next proposition we prove a partial converse of above Corollary 5.8.

Theorem 5.9. *Let I be a regular ideal. If $\text{ann } M = 0$ then $\mathcal{F}_M^I$ is a stable I -filtration.*

Proof. Notice that $M \neq H_I^0(M)$. Set $N = M/H_I^0(M)$. Note that $\text{grade}(I, N) > 0$. We have

$$(*) \quad r(I^n, M) = r(I^n, N) \quad \text{for all } n \geq 0. \quad (\text{by Lemma 3.2})$$

Let $x \in \text{ann}_R(N)$. We have $xM \subseteq H_I^0(M)$. Thus there exists $k \geq 1$ such that $I^k(xM) = 0$, so $xI^k \subseteq \text{ann } M = 0$. But since I is regular, $x = 0$. Hence $\text{ann}_R(N) = 0$. Therefore, by Theorem 5.3, the filtration $\{r(I^n, N)\}_{n \geq 0}$ is a stable I -filtration and so is $\mathcal{F}_M^I$. $\square$

6. THE CASE WHEN $M_{\mathfrak{p}}$ IS FREE FOR ALL $\mathfrak{p} \in \text{Spec}(R) \setminus \text{m-Spec}(R)$.

In this section we study our filtration $\mathcal{F}_M^I = \{r(I^n, M)\}_{n \geq 1}$ when M is free for all $\mathfrak{p} \in \text{Spec } R \setminus \text{m-Spec } R$. We show that for a regular ideal I , $\mathcal{F}_M^I$ is a stable I -filtration when $\text{Ass } R \cap \text{m-Spec } R = \emptyset$. We also prove that if $A^*(I) \cap \text{m-Spec } R = \emptyset$ then $r(I^n, M) = \widehat{I}^n$ for all $n \geq 1$. Here $A^*(I)$ is the stable value of the sequence $\text{Ass}(A/I^n)$.

6.1. Throughout this section we assume that

- (1) $\text{Ass } R \cap \text{m-Spec } R = \emptyset$ and
- (2) M is an R -module such that $M_{\mathfrak{p}}$ is free for all $\mathfrak{p} \in \text{Spec } R \setminus \text{m-Spec } R$, where $\text{m-Spec } R = \{\mathfrak{m} : \mathfrak{m} \text{ is a maximal ideal of } R\}$.

We give some examples where these assumptions hold.

Examples 6.2.

- (1) $\text{Ass } R \cap \text{m-Spec } R = \emptyset$ holds if and only if $\text{depth } R_{\mathfrak{m}} > 0$ for all $\mathfrak{m} \in \text{m-Spec } R$. Thus if $(R, \mathfrak{m})$ is a local domain which is not a field then assumption 6.1(1) holds.
- (2) If an R -module M satisfies the exact sequence of the form

$$0 \longrightarrow M \longrightarrow F \longrightarrow F/M \longrightarrow 0, \quad \text{with } \ell(F/M) < \infty,$$

where F is a free R -module then $M_{\mathfrak{p}}$ is free for all $\mathfrak{p} \in \text{Spec } R \setminus \text{m-Spec } R$.

- (3) Let $(R, \mathfrak{m})$ be a local Cohen-Macaulay ring and an isolated singularity i.e., $R_{\mathfrak{p}}$ is regular local for all prime $\mathfrak{p} \neq \mathfrak{m}$. Then if M is a maximal Cohen-Macaulay R -module then $M_{\mathfrak{p}}$ is free for all $\mathfrak{p} \neq \mathfrak{m}$.

Lemma 6.3. *(with hypotheses as in 6.1) $\text{ann } M = 0$.*

Proof. Notice that $(\text{ann } M)_{\mathfrak{p}} = \text{ann}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) = 0$, for all $\mathfrak{p} \in \text{Spec } R \setminus \text{m-Spec } R$. By our first hypothesis, we can have $\text{Ass}(\text{ann } M) \cap \text{m-Spec } R = \emptyset$ and so $\text{Ass}(\text{ann } M) = \emptyset$. Therefore $\text{ann } M = 0$. $\square$

The following proposition readily follows from Theorem 5.9 and Lemma 6.3.

Proposition 6.4. *(with hypotheses as in 6.1) If I is a regular ideal then $\mathcal{F}_M^I$ is a stable I -filtration and so $\mathcal{R}(\mathcal{F}_M^I)$ is finitely generated as an $\mathcal{R}(I)$ -module.*

Remarks 6.5.

- (1) By the result of Brodmann [2], the sequence $\text{Ass}(R/I^n)$ stabilizes for large n . Let $A^*(I)$ denote the stable value of this sequence.
- (2) Ratliff in his paper [10, 2.7], has proved that the sequence $\text{Ass}(R/\widetilde{I^n})$ eventually stabilizes at a set denoted by $\overline{A^*}(I)$.
- (3) In [10, 2.8], it is also proved that $\overline{A^*}(I) \subseteq A^*(I)$.
- (4) By [12, 1.6], we have $\mathfrak{p} \in A^*(I)$ if and only if $\mathfrak{p}_S \in A^*(I_S)$, for any multiplication closed set S disjoint from $\mathfrak{p}$.

The following is well-known. We include a proof for lack of a suitable reference.

Lemma 6.6. *For a regular ideal I we have*

- (a) $\text{Ass}(R/\widetilde{I^n}) \subseteq \text{Ass}(R/\widetilde{I^{n+1}})$ for all $n \geq 1$.
- (b) Further, $\text{Ass}(R/\widetilde{I^n}) \subseteq A^*(I)$ for all $n \geq 1$.

Proof. (a) Fix $n \geq 1$. Let $\mathfrak{p} \supseteq I$ be such that $\mathfrak{p} \in \text{Ass}(R/\widetilde{I^n})$. We localize R at $\mathfrak{p}$. Set $\mathfrak{m} = \mathfrak{p}R_{\mathfrak{p}}$. Since associated primes behave well with respect to localization so we may assume that $(R, \mathfrak{m})$ is local and $\mathfrak{m} \in \text{Ass}(R/\widetilde{I^n})$. We may further assume that $R/\mathfrak{m}$ is infinite. Otherwise we make a base change $R \longrightarrow R[X]_{\mathfrak{m}R[X]} = T$. Let $\mathfrak{n} = \mathfrak{m}T$, the extension of the maximal ideal of R in T . Notice that if E is an R -module then

$$\mathfrak{n} \in \text{Ass}_T(E \otimes_R T) \text{ if and only if } \mathfrak{m} \in \text{Ass } E.$$

Therefore we assume that $(R, \mathfrak{m})$ is local with $R/\mathfrak{m}$ infinite and $\mathfrak{m} \in \text{Ass}(R/\widetilde{I^n})$. Let $x \in I$ be a superficial element with respect to I . Consider the map

$$\mu_n^x : R/\widetilde{I^n} \longrightarrow R/\widetilde{I^{n+1}}, \quad \text{such that } a + \widetilde{I^n} \longmapsto ax + \widetilde{I^{n+1}}.$$

Clearly μ_n^x is R -linear. Also it is injective. Thus $\mathfrak{m} \in \text{Ass}(R/\widetilde{I^n}) \subseteq \text{Ass}(R/\widetilde{I^{n+1}})$.

(b) By repeatedly using (a) we get

$$\mathfrak{p} \in \text{Ass}(R/\widetilde{I^{n+k}}) \text{ for all } k \geq 1.$$

Note that for $k \gg 0$, $\widetilde{I^{n+k}} = I^{n+k}$. Also by Remark 6.5(1), $\text{Ass}(R/I^{n+k}) = A^*(I)$. Therefore the result follows. $\square$

Theorem 6.7. *(with hypotheses as in 6.1) Let I be a regular ideal of R . Then the function $n \mapsto \ell(r(I^n, M)/\widetilde{I^n})$ is a polynomial function. Furthermore if $A^*(I) \cap \text{m-Spec } R = \emptyset$ then $r(I^n, M) = \widetilde{I^n}$ for all $n \geq 1$.*

Proof. Notice that $\widetilde{I^n} \subseteq r(I^n, M)$ for all $n \geq 1$. Let $\mathfrak{p} \in \text{Spec } R \setminus \text{m-Spec } R$. By hypotheses, Proposition 1.4(e) and Remark 1.5, we have

$$(*) \quad r(I^n, M)_{\mathfrak{p}} = r_{R_{\mathfrak{p}}}(I_{\mathfrak{p}}^n, M_{\mathfrak{p}}) = \widetilde{I_{\mathfrak{p}}^n} \text{ for all } n \geq 1.$$

Therefore for all $n \geq 1$,

$$\ell\left(\frac{r(I^n, M)}{\widetilde{I^n}}\right) \text{ is finite.}$$

By Proposition 6.4, $\mathcal{R}(\mathcal{F}_M^I)$ is a finitely generated $\mathcal{R}(I)$ -module. Also $\widetilde{\mathcal{R}}(I)$ is a finitely generated $\mathcal{R}(I)$ -module. So

$$E = \frac{\mathcal{R}(\mathcal{F}_M^I)}{\widetilde{\mathcal{R}}(I)} = \bigoplus_{n \geq 1} \frac{r(I^n, M)}{\widetilde{I}^n}$$

is a finitely generated $\mathcal{R}(I)$ -module. From a well-known fact it follows that the function $n \mapsto \ell(r(I^n, M)/\widetilde{I}^n)$ is a polynomial function.

The exact sequence $0 \rightarrow r(I^n, M)/\widetilde{I}^n \rightarrow R/\widetilde{I}^n$ yields

$$\text{Ass}\left(\frac{r(I^n, M)}{\widetilde{I}^n}\right) \subseteq \text{Ass}\left(R/\widetilde{I}^n\right) \subseteq A^*(I).$$

If $A^*(I) \cap \text{m-Spec } R = \emptyset$ then by (*) we get $r(I^n, M) = \widetilde{I}^n$ for all $n \geq 1$. $\square$

Remark 6.8. If $A^*(I) \cap \text{m-Spec } R \neq \emptyset$ then $r(I, M)$ need not be equal to $\widetilde{I}$. For instance let $R = k[t^4, t^{11}, t^{17}, t^{18}]_{\mathfrak{m}}$ where $\mathfrak{m} = (t^4, t^{11}, t^{17}, t^{18})$. Let $I = (t^4, t^{11})$. Set $M = (t^4, t^{11}, t^{17})$ considered as a submodule of R . Notice that $\ell(R/M)$ is finite. By Example 9.7 we get that $r(I^n, M) \neq \widetilde{I}^n$ for all $n \geq 1$. In this case I is $\mathfrak{m}$ -primary. So $A^*(I) = \{\mathfrak{m}\}$.

In view of Theorem 6.7 we give some situations of prime $\mathfrak{p}$ such that $\mathfrak{p} \in A^*(I)$.

Remarks 6.9.

- (1) If $ht(\mathfrak{p}) = l(I_{\mathfrak{p}})$, the *analytic spread* of $I_{\mathfrak{p}}$ then $\mathfrak{p} \in A^*(I)$. (see [12, 4.1])
- (2) If $\mathfrak{p} \in A^*(I) \setminus B^*(I)$, where $B^*(I)$ is the stable value of the sequence $\text{Ass}(I^n/I^{n+1})$, then $\mathfrak{p} \in \text{Ass } R$. (see [12, 2.2])
- (3) With hypotheses as in 6.1, if a maximal ideal $\mathfrak{m} \in A^*(I)$ then $\mathfrak{m} \in B^*(I)$ that is, $\mathfrak{m} \in \text{Ass}_R(I^n/I^{n+1})$ for $n \gg 0$. So we get $\mathfrak{m}/I \in \text{Ass}_{R/I}(I^n/I^{n+1})$ for $n \gg 0$. Set $G = G_I(R)$. By [13, 2.1], we thus have $\mathfrak{m}/I = Q \cap R/I$ such that $Q \in \text{Ass}(G) \setminus V(G_+)$.

Corollary 6.10. (with hypotheses as in 6.1) In addition let $(R, \mathfrak{m})$ be a local Cohen-Macaulay ring. Let $x_1, \dots, x_r$ be a regular sequence with $r < \dim R$. Set $I = \langle x_1, \dots, x_r \rangle R$. Then $r(I^n, M) = I^n$ for all $n \geq 1$.

Proof. Clearly R/I^n is a Cohen-Macaulay ring of dimension greater than or equal to 1 for all $n \geq 1$. So $\mathfrak{m} \notin \text{Ass}(R/I^n)$ for all $n \geq 1$. This gives $\mathfrak{m} \notin A^*(I)$. Therefore by Theorem 6.7, $r(I^n, M) = \widetilde{I}^n$ for all $n \geq 1$. However as $\text{depth } G_I(R) > 0$ we get $\widetilde{I}^n = I^n$ for all $n \geq 1$. The result follows. $\square$

7. SOME MORE ANALYSIS ON $r(I, M)$

In this section we analyze the case when $\text{ann } M$ need not be zero. We also consider the case when $\text{grade}(I, M) = 0$. When $M \neq H_I^0(M)$ both these cases can be dealt with by going modulo the ideal $(H_I^0(M) : M)$. We prove that $r(I^{n+1}, M) = I \cdot r(I^n, M) + (H_I^0(M) : M)$ for all $n \gg 0$. Our techniques also yield $\widetilde{I}^n M = I^n M + H_I^0(M)$ for all $n \gg 0$.

7.1. Before we proceed further let us fix some notations which we will use throughout the section. Set $N = M/H_I^0(M)$, $q_I(M) = (H_I^0(M) : M)$ and $S = R/q_I(M)$. Let J be the image of I in S .

Proposition 7.2. *Let M be an R -module such that $M \neq H_I^0(M)$. Then*

- (a) $r(I^n, M) = r_R(I^n, N)$ for all $n \geq 1$.
- (b) $q_I(M) \subseteq r(I^n, N)$ for all $n \geq 1$.
- (c) $\text{ann}_S(N) = 0$.
- (d) $\text{grade}_R(I, N) = \text{grade}_S(J, N) > 0$.
- (e) $r(I^n, M)/q_I(M) = r_S(J^n, N)$ for all $n \geq 1$.

Proof. Set $q = q_I(M)$. (a) follows from Lemma 3.2(a). Parts (b) and (c) are easy to prove. For (d), note that $H_J^0(N) = 0$. To prove (e), it is sufficient to show that

$$\frac{r(I^n, N)}{q} = r_S(J^n, N) \quad \text{for all } n \geq 1.$$

Let $x \in r(I^n, N)$. Thus $xI^k N \subseteq I^{n+k}N$, so by going modulo q , we get

$$\bar{x}J^k N \subseteq J^{n+k}N. \text{ So } \bar{x} \in r_S(J^n, N).$$

Conversely if $\bar{x} \in r_S(J^n, N)$ then we have $\bar{x}J^k N \subseteq J^{n+k}N$. Thus

$$\begin{aligned} (x+q) \left(\frac{I^k + q}{q} \right) N &\subseteq \left(\frac{I^{n+k} + q}{q} \right) N, \\ \text{so } \left(\frac{xI^k + q}{q} \right) N &\subseteq \left(\frac{I^{n+k} + q}{q} \right) N. \end{aligned}$$

This implies $(xI^k + q)N \subseteq (I^{n+k} + q)N$. So $xI^k N \subseteq I^{n+k}N$, since $q = \text{ann}_R N$. Therefore $x \in r(I^n, N)$ and hence $\bar{x} \in r(I^n, N)/q$. $\square$

Theorem 7.3. *Let M be an R -module such that $M \neq H_I^0(M)$. Then*

$$r(I^{n+1}, M) = I \cdot r(I^n, M) + (H_I^0(M) : M) \quad \text{for all } n \gg 0.$$

Proof. By Proposition 7.2(d), $\text{grade}_S(J, N) > 0$. Together with result (c), this gives that the filtration $\{r_S(J^n, N)\}_{n \geq 0}$ is a stable J -filtration (by Theorem 5.3). Therefore we have $r_S(J^{n+1}, N) = J \cdot r_S(J^n, N)$ for all $n \gg 0$. So

$$\frac{r(I^{n+1}, M)}{q} = \left(\frac{I + q}{q} \right) \cdot \left(\frac{r(I^n, M)}{q} \right).$$

Thus $r(I^{n+1}, M) = I \cdot r(I^n, M) + q$ for all $n \gg 0$. $\square$

7.4. Consequences of Theorem 7.3

- (1) When $M = R$ we have, for any ideal I ,

$$\widetilde{I^{n+1}} = I \cdot \widetilde{I^n} + H_I^0(R) \quad \text{for all } n \gg 0.$$

- (2) If $\text{grade}(I, M) > 0$ then

$$r(I^{n+1}, M) = I \cdot r(I^n, M) + \text{ann } M \quad \text{for all } n \gg 0.$$

Next we relate $\widetilde{I^n M}$ and $I^n M$. In the case when $M = R$, the following result is proved in [21, 2.13(a)].

Proposition 7.5. *Let I be an ideal of R and M an R -module. Then*

$$\widetilde{I^n M} = I^n M + H_I^0(M) \quad \text{for all } n \gg 0.$$

Proof. By Proposition 7.2(d), $\text{grade}(J, N) > 0$. Therefore $\widetilde{J^n N} = J^n N$ for all $n \gg 0$. So

$$(i) \quad \widetilde{J^n N} = \frac{I^n M + H_I^0(M)}{H_I^0(M)} \quad \text{for all } n \gg 0.$$

It is easy to see that $H_I^0(M) \subseteq \widetilde{I^n M}$ for all $n \geq 1$. By an argument similar to Proposition 7.2(e), we get

$$\widetilde{J^n N} = \frac{\widetilde{I^n M}}{H_I^0(M)}.$$

Thus from equation (i), the result follows. $\square$

Corollary 7.6. *Assume that $\widetilde{\mathcal{R}(I, M)} = \bigoplus_{n \geq 0} \widetilde{I^n M}$ is a Noetherian $\mathcal{R}(I)$ -module and M is separated with respect to the I -adic topology. Then $H_I^0(M) = 0$.*

Proof. Since $\widetilde{\mathcal{R}(I, M)}$ is Noetherian, there exists a positive integer $n_0 \in \mathbb{N}$ such that $\widetilde{I^n M} = I^{n-n_0} \widetilde{I^{n_0} M}$ for all $n \geq n_0$. So $H_I^0(M) \subseteq I^{n-n_0} \widetilde{I^{n_0} M} \subseteq I^{n-n_0} M$ for all $n \geq n_0$. Thus the result follows from our hypothesis on M . $\square$

8. RELATION WITH A SUPERFICIAL ELEMENT

In this section we assume that $(R, \mathfrak{m})$ is local with the maximal ideal $\mathfrak{m}$. The goal of this section is to understand the relation between $r(I, M)$ and a superficial element. To ensure the existence of superficial element we assume (unless stated otherwise) that the residue field $K = R/\mathfrak{m}$ is infinite. When I is $\mathfrak{m}$ -primary our techniques yield an easily computable bound on k such that $\widetilde{I^n} = (I^{n+k} : I^k)$ for all $n \geq 1$.

8.1. Recall an element $x \in I$ is called *M-superficial* with respect to I if there exists an integer $c \geq 0$ such that

$$(I^{n+1}M :_M x) \cap I^c M = I^n M \quad \text{for all } n \geq c.$$

Superficial element exists when K is infinite. If $\text{grade}(I, M) > 0$ then every M -superficial element is also M -regular. Also if $x \in I$ is M -superficial and M -regular then

$$(I^{n+1}M :_M x) = I^n M \quad \text{for all } n \gg 0. \quad (\text{see [20, p. 8] for the case } M = R)$$

Proposition 8.2. *Let $x \in I$ be a M -superficial element. Then*

$$(r(I^{n+1}, M) : x) = r(I^n, M) \quad \text{for all } n \geq 1.$$

Proof. Since $x \in I$ is M -superficial, there exists $c > 0$ such that

$$(i) \quad (I^{j+1}M :_M x) \cap I^c M = I^j M \quad \text{for all } j \geq c.$$

It is easy to see that $r(I^n, M) \subseteq (r(I^{n+1}, M) : x)$ for all $n \geq 1$. Conversely let $a \in (r(I^{n+1}, M) : x)$. Then

$$ax \in r(I^{n+1}, M) = (I^{n+k+1}M : I^k M) \quad \text{for } k \gg 0$$

$$\text{Thus } axI^k M \subseteq I^{n+k+1}M \quad \text{for } k \gg 0.$$

Now for $k \gg 0$, we have $aI^k M \subseteq I^c M$. Therefore, by (i), we get

$$aI^k M \subseteq I^{n+k}M.$$

Thus $a \in (I^{n+k}M : I^k M) \subseteq r(I^n, M)$. $\square$

8.3. If $\text{grade}(I, M) > 0$ then $\widetilde{I^n M} = I^n M$ for all $n \gg 0$. Define

$$\rho^I(M) := \min\{ n : \widetilde{I^i M} = I^i M \text{ for all } i \geq n \}.$$

If $x \in I$ is M -superficial then it is proved in [16, Corollary 5.3] that

$$\rho^I(M) = \min\{ n : (I^{i+1}M :_M x) = I^i M \text{ for all } i \geq n \}.$$

Proposition 8.4. *Let $\text{grade}(I, M) > 0$. Then*

$$r(I^n, M) = (I^n M : M) \quad \text{for all } n \geq \rho^I(M).$$

Proof: Let $x \in I$ be both M -superficial and M -regular. So

$$(i) \quad (I^{n+1}M :_M x) = I^n M \quad \text{for all } n \geq \rho^I(M).$$

Let $a \in r(I^n, M)$. Then $aI^k M \subseteq I^{n+k}M$ for some $k \geq 1$. So $ax^k M \subseteq I^{n+k}M$. By repeated use of (i), we get $aM \subseteq (I^{n+k}M :_M x^k) = I^n M$. Therefore $a \in (I^n M : M)$. $\square$

Corollary 8.5. *Let $x \in I$ be M -superficial such that x^* is $G_I(M)$ -regular. Then*

$$r(I^n, M) = (I^n M : M) \quad \text{for all } n \geq 1.$$

Proof. Since x^* is $G_I(M)$ -regular, we have

$$(I^{n+1}M :_M x) = I^n M \quad \text{for all } n \geq 1$$

So $\rho^I(M) = 1$. Therefore the result follows from Proposition 8.4. $\square$

Notation: Let I be an $\mathfrak{m}$ -primary ideal. Suppose M is of dimension $d \geq 0$. We define the postulation number $\eta^I(M)$ of I with respect to M as follows:

$$\eta^I(M) := \min\{ n : H_I^M(t) = p_I^M(t), \text{ for all } t \geq n \},$$

where $H_I^M(n) = \ell(M/I^{n+1}M)$ is the Hilbert-Samuel function of M with respect to I and $p_I^M(t)$ is the Hilbert-Samuel polynomial of M with respect to I . Let $x \in I$ be M -superficial. We set

$$\eta^I(x, M) = \max\{ \eta^I(M), \eta^I(M/xM) \},$$

The following proposition is proved by J. Elias in [6, 1.3] for $M = R$. The same proof applies to the general case.

Proposition 8.6. *Let I be an $\mathfrak{m}$ -primary ideal. Let $x \in I$ be an M -superficial and M -regular element. Then*

$$(I^{k+1}M : x) = I^k M \quad \text{for all } k \geq \eta^I(x, M) + 1.$$

$\square$

Remark 8.7. From above Proposition 8.6, if $\text{grade}(I, M) > 0$, it follows that $\rho^I(M) \leq \eta^I(x, M) + 1$.

Proposition 8.8. *Let $\text{grade}(I, M) > 0$. Then*

$$r(I^n, M) = (I^{n+k}M : I^k M) \quad \text{for all } n \geq 1 \text{ and } k \geq \rho^I(M).$$

Proof. Clearly $\text{grade}(I, I^k M) > 0$ for all k . By Proposition 1.4(g) and Proposition 2.7, we have for each $n \geq 1$,

$$\begin{aligned} r(I^n, M) &= r(I^n, I^k M) \quad \text{for all } k \geq 1, \\ &= (\widetilde{I^{n+k} M} : I^k M) \quad \text{for all } k \geq 1. \\ \text{Thus } r(I^n, M) &= (I^{n+k} M : I^k M) \quad \text{for all } k \geq \rho^I(M). \end{aligned}$$

□

When $M = R$ we obtain that

Corollary 8.9. *Let I be a regular ideal. Then for each value of n , we have*

$$\widetilde{I^n} = (I^{n+k} : I^k) \quad \text{for all } k \geq \rho^I(R).$$

Remark 8.10. In particular, if I is an $\mathfrak{m}$ -primary regular ideal then for each $n \geq 1$,

$$\widetilde{I^n} = (I^{n+k} : I^k) \quad \text{for all } k \geq \eta^I(x, R) + 1.$$

For $n = 1$ the result above was proved by Elias [6, p. 722]. However our result does not follow from it. Furthermore even for $n = 1$ our method is simpler to compute.

It is of interest to find a similar bound for $\widetilde{I^n M}$. We prove

Theorem 8.11. *Let $\text{grade}(I, M) > 0$. Then for each $n \geq 1$, we have*

$$\widetilde{I^n M} = (I^{n+k} M :_M I^k) \quad \text{for all } k \geq \rho^I(M).$$

Proof. Let $x \in I$ be M -superficial. It is enough to show that for $k \geq \rho^I(M)$, we have

$$(I^{n+k+1} M :_M I^{k+1}) \subseteq (I^{n+k} M :_M I^k).$$

Let $m \in (I^{n+k+1} M :_M I^{k+1})$. Then $m I^{k+1} \subseteq I^{n+k+1} M$, so $m x I^k \subseteq I^{n+k+1} M$. Therefore

$$m I^k \subseteq (I^{n+k+1} M :_M x) = I^{n+k} M.$$

So $m \in (I^{n+k} M :_M I^k)$. □

The next theorem deals with the situation when residue field $R/\mathfrak{m}$ is not necessarily infinite.

Theorem 8.12. *Let $(R, \mathfrak{m})$ be local, M an R -module and let I be an ideal with $\text{grade}(I, M) > 0$. Then the following hold for all $k \geq \rho^I(M)$,*

- (a) $r(\widetilde{I^n}, M) = (I^{n+k} M : I^k M)$ for all $n \geq 1$.
- (b) $\widetilde{I^n M} = (I^{n+k} M :_M I^k)$ for all $n \geq 1$.

In particular, when $M = R$ we have for each n , $\widetilde{I^n} = (I^{n+k} : I^k)$ for all $k \geq \rho^I(R)$.

Proof. Note that (b) follows from (a). We now prove part (a). Consider the faithfully flat extension

$$R \longrightarrow T = R[X]_{\mathfrak{m}R[X]}.$$

Note that the residue field of T is $K(X)$, the quotient field of polynomial ring $K[X]$ and it is infinite. Set $q = IT$ and $E = M \otimes_R T$. By Proposition 1.4(d), we get

$$\begin{aligned} r(I^n, M) \otimes_R T &= r(q^n, E) \\ &= (q^{n+k} E :_T q^k E) \quad \text{for all } k \geq \rho^q(E), \\ &= (I^{n+k} M : I^k M) \otimes_R T \quad \text{for all } k \geq \rho^q(E). \end{aligned}$$

By [16, 1.7], we have $\rho^q(E) = \rho^I(M)$. Fix $k \geq \rho^I(M)$ and set

$$D = r(I^n, M) / (I^{n+k} M : I^k M).$$

Then we have $D \otimes_R T = 0$. Since T is a faithfully flat extension of R , we get $D = 0$. $\square$

We used the packages *CoCoA* [1] and *Singular* [7] for our computations. We reconsider the Example 1.3. In this example we apply Theorem 8.12 to compute $\widetilde{I}^n$ for each n .

Example 8.13. (see [17, 1.4]) Let $R = k[x, y]_{\langle x, y \rangle}$ and the ideal

$$I = \langle y^{22}, x^4 y^{18}, x^7 y^{15}, x^8 y^{14}, x^{11} y^{11}, x^{14} y^8, x^{15} y^7, x^{18} y^4, x^{22} \rangle R.$$

Set $u = x^{22} + y^{22}$. The Poincare series of I and I/u are

$$\begin{aligned} PS_I(t) &= \frac{227 + 189t + 10t^2 + 10t^3 - 2t^4}{(1-t)^2}, \\ PS_{I/u}(t) &= \frac{227 + 189t + 12t^2 + 6t^3}{(1-t)}. \end{aligned}$$

So $e_i(I/u) = e_i(I)$ for $i = 0, 1$. Thus u is R -superficial with respect to I . Note that $\rho^I(R) \leq \eta^I(x, R) = 3$. Therefore by 8.12, $\widetilde{I}^n = (I^{n+3} : I^3) = I^n$ for all $n \geq 3$. Also

$$\begin{aligned} \widetilde{I} &= (I^4 : I^3) = I, \\ \widetilde{I}^2 &= (I^5 : I^3) = I^2 + \langle x^{20} y^{24}, x^{24} y^{20} \rangle R. \end{aligned}$$

9. THE CASE WHEN I HAS A PRINCIPAL REDUCTION

In this section we discuss ideals having principal reductions. When I has a principal reduction $J = (x)$, the computation of $r(I, M)$ is greatly simplified. Let $r = r_x(I) = \min\{n : I^{n+1} = xI^n\}$, the reduction number of I with respect to (x) then we show that $r(I^n, M) = (I^{n+r} M : I^r M)$ for all $n \geq 1$. This result is then used to compute many examples.

Proposition 9.1. *Let $\text{grade}(I, M) > 0$. If I has a principal reduction (x) with reduction number $r_x(I) \leq r$ then the following hold*

- (a) *If $r(I^{n+1}, M) = x \cdot r(I^n, M)$ for some n then $r(I^{m+1}, M) = x \cdot r(I^m, M)$ for all $m \geq n$.*
- (b) *$r(\widetilde{I^n}, M) = (I^{n+r} M : I^r M)$ for all $n \geq 1$.*
- (c) *$\widetilde{I^n M} = (I^{n+r} M :_M I^r)$ for all $n \geq 1$.*

But before we do this we first need to prove the following lemma.

Lemma 9.2. *Let the situation be as in Proposition 9.1. Then x is M -regular and*

$$(x) \cap r(I^{n+1}, M) = x \cdot r(I^n, M) \text{ for all } n \geq 1.$$

Proof. It is easy to check that x is M -regular. Let $ax \in r(I^{n+1}, M)$ for some $a \in R$. Then $axI^k M \subseteq I^{n+1+k} M$ for some $k \geq 0$. We assume $k \geq r_x(I)$. Therefore

$$axI^k M \subseteq xI^{n+k} M.$$

Since x is M -regular, we get $a \in (I^{n+k} M : I^k M) \subseteq r(I^n, M)$. $\square$

We now give the proof of proposition.

Proof of Proposition 9.1. (a) By Lemma 9.2, x is M -regular. For all $m \geq n$, we have

$$\begin{aligned} r(I^{m+1}, M) &\subseteq r(I^{n+1}, M) \subseteq (x). \\ \text{Thus, } r(I^{m+1}, M) &\subseteq (x) \cap r(I^{m+1}, M), \\ &= x \cdot r(I^m, M), \quad (\text{by Lemma 9.2}) \\ &\subseteq r(I^{m+1}, M). \end{aligned}$$

(b): It is enough to show that $(I^{n+r+1} M : I^{r+1} M) \subseteq (I^{n+r} M : I^r M)$. Let $a \in (I^{n+r+1} M : I^{r+1} M)$.

$$\begin{aligned} aI^{r+1} M &\subseteq I^{n+r+1} M, \\ \text{so } ax \cdot I^r M &\subseteq x \cdot I^{n+r} M. \end{aligned}$$

As x is M -regular we get $aI^r M \subseteq I^{n+r} M$.

(c) It is similar to (b). $\square$

For $M = R$ Proposition 9.1(c) yields

Corollary 9.3. *Let I be a regular ideal having a principal reduction (x) with reduction number $r_x(I) \leq r$. Then $\widetilde{I}^n = (I^{n+r} : I^r)$ for all $n \geq 1$. $\square$*

Remark 9.4. For $n = 1$ the result above has been proved already in [4, 2.1]. However for $n \geq 2$ their result does not imply ours. Notice that, since $r_{x^n}(I^n) \leq r_x(I) \leq r$, [4, 2.1] implies $\widetilde{I}^n = (I^{nr+n} : I^{nr})$.

Examples: We use Proposition 9.1 to construct many examples. We first give an example of a module M with no free summand such that $r(I^n, M) = I^n$ for all $n \geq 1$ but $IM \neq \widetilde{IM}$.

Example 9.5. Let $Q = k[x, y]_{\langle x, y \rangle}$ be a local ring with the maximal ideal $\mathfrak{n}$. Set $(R, \mathfrak{m}) = (Q/\langle y^3 \rangle, \mathfrak{n}/\langle y^3 \rangle)$. Consider the R -module

$$M = \left\langle \begin{pmatrix} 0 \\ y^2 \end{pmatrix}, \begin{pmatrix} y \\ x \end{pmatrix} \right\rangle \subseteq R^2.$$

Note that M is a Cohen-Macaulay R -module. Also notice that $\mathfrak{m}^3 = x \cdot \mathfrak{m}^2$ so $r_x(\mathfrak{m}) = 2$. We compute $r(\mathfrak{m}^n, M)$ for $n \geq 1$. Using Proposition 9.1(b), one checks

$$r(\mathfrak{m}^i, M) = (\mathfrak{m}^{i+2} M : \mathfrak{m}^2 M) = \mathfrak{m}^i \text{ for } i = 1, 2, 3.$$

Thus, by Proposition 9.1(a) we have

$$r(\mathfrak{m}^{n+1}, M) = x \cdot r(\mathfrak{m}^n, M) = \mathfrak{m}^{n+1} \text{ for all } n \geq 2.$$

We also compute $\widetilde{\mathfrak{m}M}$ by using Proposition 9.1(c)

$$\begin{aligned}\widetilde{\mathfrak{m}M} &= (\mathfrak{m}^3 M :_M \mathfrak{m}^2) = \left\langle \begin{pmatrix} 0 \\ y^2 \end{pmatrix}, \begin{pmatrix} y^2 \\ xy \end{pmatrix}, \begin{pmatrix} xy \\ x^2 \end{pmatrix} \right\rangle \\ \text{and } \mathfrak{m}M &= \left\langle \begin{pmatrix} y^2 \\ xy \end{pmatrix}, \begin{pmatrix} xy \\ x^2 \end{pmatrix} \right\rangle.\end{aligned}$$

Hence $\mathfrak{m}M \neq \widetilde{\mathfrak{m}M}$.

We give an example where $r(I^n, M) = \widetilde{I}^n$ for all $n \geq 1$ and $IM = \widetilde{IM}$.

Example 9.6. Let

$$A = \frac{k[x, y, z]}{\langle x^5 - z^2, y^3 - xz \rangle} \simeq k[t^6, t^7, t^{15}].$$

Set $R = A_{\mathfrak{n}}$, where $\mathfrak{n} = \langle x, y, z \rangle A$. Let $\mathfrak{m}$ be the maximal ideal of R . Let

$$M = \left\langle \begin{pmatrix} x^2 \\ z^2 \end{pmatrix}, \begin{pmatrix} y \\ yz \end{pmatrix} \right\rangle \subseteq R^2.$$

Note that M is a Cohen-Macaulay R -module and $\mathfrak{m}^6 = x \cdot \mathfrak{m}^5$ so $r_x(\mathfrak{m}) = 5$. Therefore using Proposition 9.1(b), one can check the following

$$r(\mathfrak{m}^n, M) = (\mathfrak{m}^{n+5} M : \mathfrak{m}^5 M) = \widetilde{\mathfrak{m}}^n = (\mathfrak{m}^{n+5} : \mathfrak{m}^5) \text{ for } n = 1, \dots, 6.$$

$$\text{Also } \widetilde{\mathfrak{m}}^n \neq \mathfrak{m}^n \text{ for } n = 2, 3, 4 \text{ and } \widetilde{\mathfrak{m}}^n = \mathfrak{m}^n \text{ for all } n \geq 5.$$

Thus, by Proposition 9.1(a) we have

$$r(\mathfrak{m}^{n+1}, M) = x \cdot r(\mathfrak{m}^n, M) = \mathfrak{m}^{n+1} \text{ for all } n \geq 5.$$

Using Proposition 9.1(c), one can verify that $\widetilde{\mathfrak{m}M} = (\mathfrak{m}^6 M :_M \mathfrak{m}^5) = \mathfrak{m}M$.

Next we give an example in which we have $\widetilde{I}^n \neq r(I^n, M)$ for all $n \geq 1$, and $\mathcal{F}_M^I$ is a stable I -filtration.

Example 9.7. Let $A = k[t^4, t^{11}, t^{17}, t^{18}]$. Using Singular we get

$$A \simeq B = \frac{k[x, y, z, w]}{\langle y^2 - xw, yz - x^7, z^2 - x^4w, yw - x^3z, zw - x^6y, w^2 - x^2yz \rangle}.$$

Set $R = B_{\mathfrak{m}}$, where $\mathfrak{m} = \langle x, y, z, w \rangle B$. Let $I = \langle x, y \rangle R$ and $M = \langle x, y, z \rangle R$. One can check $r_x(I) = 2$. Note that x is both M -regular and R -regular. By Proposition 9.1(b), we have

$$\begin{aligned}r(I, M) &= (I^3 M : I^2 M) = \langle x, y, z, w \rangle R, \\ r(I^2, M) &= (I^4 M : I^2 M) = \langle x^2, xy, xz, xw \rangle R = x \cdot r(I, M). \\ \text{And } \widetilde{I} &= (I^3 : I^2) = \langle x, y, w \rangle R, \\ \widetilde{I}^2 &= (I^4 : I^2) = \langle x^2, xy, xw \rangle R = x \cdot \widetilde{I}.\end{aligned}$$

Therefore we have

$$\begin{aligned}r(I^{n+1}, M) &= x \cdot r(I^n, M) = \langle x^{n+1}, x^n y, x^n z, x^n w \rangle R \text{ for all } n \geq 1, \\ \widetilde{I^{n+1}} &= x \cdot \widetilde{I}^n = \langle x^{n+1}, x^n y, x^n w \rangle R \text{ for all } n \geq 1.\end{aligned}$$

Notice that $x^{n-1}z \leftrightarrow t^{4n+13}$. We use the expression of I^n from Example 1.2 to verify that $x^{n-1}z \notin \widetilde{I}^n$. Therefore $\widetilde{I}^n \neq r(I^n, M)$ for all $n \geq 1$. Since M is a

non-zero ideal of R and R is a domain so $\text{ann } M = 0$. Notice that I is a regular ideal. So, by Theorem 5.9, $\mathcal{F}_M^I$ is a stable I -filtration.

10. APPLICATION TO HILBERT FUNCTIONS

In this section we assume that $(R, \mathfrak{m})$ is a local ring with the maximal ideal $\mathfrak{m}$ and I an $\mathfrak{m}$ -primary ideal. Let $P_I^M(t)$ be the Hilbert-Samuel function of M with respect to I . We first show that if $\text{grade}(I, M) > 0$ then the set

$$\mathcal{H}(I) = \{ J : J \text{ is an ideal of } R \text{ such that } J \supseteq I \text{ and } P_J^M(t) = P_I^M(t) \}$$

has $r(I, M)$ as the unique maximal element. If $\dim M = 1$ and $\text{depth } M = 0$ then $\widetilde{\mathcal{R}}(I, M) = \bigoplus_{n \geq 0} \widetilde{I^n M}$ is not Noetherian. However $\widetilde{G}_I(M) = \bigoplus_{n \geq 0} \widetilde{I^n} / \widetilde{I^{n+1}}$ is a Noetherian $G_I(R)$ -module, in fact a Cohen-Macaulay $G_I(R)$ -module (Theorem 7.6). Next we give an application of Proposition 7.5 to show that if $\dim M = 1$ then $e_1^I(M) - e_0^I(M) + \ell(M/IM) \geq -\ell(H_{\mathfrak{m}}^0(M))$.

10.1. Recall that the Hilbert-Samuel function of M with respect to I is the function

$$n \longmapsto \ell(M/I^{n+1}M) \quad \text{for all } n \gg 0.$$

It is well known that for all large values of n it is given by a polynomial $P_I^M(n)$ of degree $r = \dim M$, the *Hilbert-Samuel polynomial* with respect to I . It can be written in the form

$$P_I^M(X) = \sum_{i=0}^r (-1)^i e_i^I(M) \binom{X+r-i}{r-i}.$$

The integers $e_0^I(M), e_1^I(M), \dots, e_r^I(M)$ are called the *Hilbert coefficients* of M with respect to I . The number $e_0^I(M)$ is also called the *multiplicity* of M with respect to I .

10.2. For an $\mathfrak{m}$ -primary ideal I , we define the set

$$\mathcal{H}(I) := \{ J : J \text{ is an ideal of } R \text{ such that } J \supseteq I \text{ and } P_J^M(t) = P_I^M(t) \}.$$

Proposition 10.3. *Let $\text{grade}(I, M) > 0$. Then $r(I, M)$ is the unique maximal element of $\mathcal{H}(I)$.*

Proof. By Corollary 2.6, we have $r(I, M)^n M = I^n M$ for all $n \gg 0$. So $r(I, M) \in \mathcal{H}(I)$. Conversely if $J \in \mathcal{H}(I)$ then $J^n M = I^n M$ for all $n \gg 0$. Let $x \in J$. We have

$$\begin{aligned} xI^{n-1}M = xJ^{n-1}M &\subseteq J^n M = I^n M \text{ for all } n \gg 0. \\ \text{So } x &\in (I^n M : I^{n-1}M) = r(I, M). \end{aligned}$$

Thus $J \subseteq r(I, M)$. It follows that $r(I, M)$ is the unique maximal element in $\mathcal{H}(I)$. $\square$

Remarks 10.4. We have

- (a) $\widetilde{I^n M} = I^n M + H_{\mathfrak{m}}^0(M)$ for all $n \gg 0$. (by Proposition 7.5)
- (b) By using Artin-Rees lemma, we can check that

$$I^n M \cap H_{\mathfrak{m}}^0(M) = 0 \quad \text{for all } n \gg 0.$$

(c) Using (a) and (b), it is easy to prove that

$$\frac{\widetilde{I^n M}}{I^n M} \simeq \frac{H_I^0(M)}{I^n M \cap H_I^0(M)} \simeq H_I^0(M) \quad \text{for all } n \gg 0.$$

Set $\widetilde{G}_I(M) = \bigoplus_{n \geq 0} \widetilde{I^n M} / \widetilde{I^{n+1} M}$. In dimension one we have

Theorem 10.5. *Let $(R, \mathfrak{m})$ be a local ring. Let I be an $\mathfrak{m}$ -primary ideal and suppose that M is an R -module with $\dim M = 1$. Then*

- (a) $\ell(M/I^{n+1}M) = e_0^I(M)(n+1) - e_1^I(M) - \ell(H_{\mathfrak{m}}^0(M))$.
- (b) $\widetilde{G}_I(M)$ is a finitely generated $G_I(R)$ -module of dimension 1.
- (c) $\widetilde{G}_I(M)$ is a Cohen-Macaulay $G_I(R)$ -module.

Proof. We may assume that $K = R/\mathfrak{m}$ is infinite, otherwise we consider the standard base change $R \rightarrow R[X]_{\mathfrak{m}_A[X]}$ (see [16, 1.3]).

(a) From the exact sequence

$$0 \rightarrow \widetilde{I^{n+1}M} / I^{n+1}M \rightarrow M / I^{n+1}M \rightarrow M / \widetilde{I^{n+1}M} \rightarrow 0$$

and using Remark 10.4(c), it follows that for all $n \gg 0$,

$$\ell(M / \widetilde{I^{n+1}M}) = e_0^I(M)(n+1) - e_1^I(M) - \ell(H_{\mathfrak{m}}^0(M)).$$

(b) Consider the exact sequence

$$0 \rightarrow \widetilde{I^n M} / \widetilde{I^{n+1} M} \rightarrow M / \widetilde{I^{n+1} M} \rightarrow M / \widetilde{I^n M} \rightarrow 0.$$

By (a), we have

$$(ii) \quad \ell(\widetilde{I^n M} / \widetilde{I^{n+1} M}) = e_0^I(M), \quad \text{for all } n \gg 0.$$

Let $x \in I$ be M -superficial which exists as the residue field K is infinite. Let x^* be the image of x in I/I^2 , considered as $G_I(R)$ -element. Then by Proposition 8.2, the sequence

$$(iii) \quad 0 \rightarrow \widetilde{I^n M} / \widetilde{I^{n+1} M} \xrightarrow{x^*} \widetilde{I^{n+1} M} / \widetilde{I^{n+2} M}$$

is exact. In view of (ii), we get

$$(iv) \quad \widetilde{I^{n+1} M} / \widetilde{I^{n+2} M} \simeq x^* \cdot (\widetilde{I^n M} / \widetilde{I^{n+1} M}) \quad \text{for all } n \gg 0, \text{ say for } n \geq n_0.$$

Let $S = \{z_1, z_2, \dots, z_s\}$ be the set of homogeneous generators of $\bigoplus_{i=0}^{n_0} \widetilde{I^n M} / \widetilde{I^{n+1} M}$ as an R -module. Using (iv) it follows that S generates $\widetilde{G}_I(M)$ as a $G_I(R)$ -module. Since $\ell(\widetilde{I^n M} / \widetilde{I^{n+1} M}) = e_0^I(M) > 0$ for all $n \gg 0$. It follows that $\dim \widetilde{G}_I(M) = 1$.

(c) By (iii), we have x^* is $\widetilde{G}_I(M)$ -regular. So $\text{depth } \widetilde{G}_I(M) > 0$. As $\dim \widetilde{G}_I(M) = 1$ we have $\widetilde{G}_I(M)$ is Cohen-Macaulay. $\square$

Theorem 10.6. *Let the situation be as in Theorem 10.5. Then*

$$e_1^I(M) - e_0^I(M) + \ell(M/IM) \geq -\ell(H_{\mathfrak{m}}^0(M)).$$

Further equality holds if and only if $\widetilde{IM} = IM$ and $\widetilde{I^n M} = I^n M + H_{\mathfrak{m}}^0(M)$ for $n \geq 2$.

In [8, 3.1] a sharper bound is found. However the case of equality is different.

Proof. As $\widetilde{G}_I(M)$ is a finitely generated $G_I(R)$ -module so there exists $h(t) \in \mathbb{Z}[t]$ such that

$$\sum_{n=0}^{\infty} \ell \left(\widetilde{I^n M} / \widetilde{I^{n+1} M} \right) t^n = \frac{h(t)}{(1-t)},$$

where $h(t) = h_0 + h_1 t + h_2 t^2 + \dots + h_s t^s$. As $\widetilde{G}_I(M)$ is a Cohen-Macaulay $G_I(R)$ -module, all the coefficients of $h(t)$ are non-negative. Thus

$$\begin{aligned} \widetilde{e}_1^I(M) &= \widetilde{e}_0^I(M) - h_0 + \sum_{j=2}^s (j-1)h_j, \\ &= \widetilde{e}_0^I(M) - \ell \left(M / \widetilde{IM} \right) + \sum_{j=2}^s (j-1)h_j, \\ &= \widetilde{e}_0^I(M) - \ell(M/IM) + \ell \left(\widetilde{IM} / IM \right) + \sum_{j=2}^s (j-1)h_j. \end{aligned}$$

For $n \gg 0$ we have $\ell \left(M / \widetilde{I^{n+1} M} \right) = \widetilde{e}_0^I(M)(n+1) - \widetilde{e}_1^I(M)$. By comparing it with 10.5(a), we get $\widetilde{e}_0^I(M) = e_0^I(M)$ and $\widetilde{e}_1^I(M) = e_1^I(M) + \ell(H_{\mathfrak{m}}^0(M))$. Thus we get

$$\begin{aligned} e_1^I(M) &= \widetilde{e}_1^I(M) - \ell(H_{\mathfrak{m}}^0(M)), \\ &= \widetilde{e}_0^I(M) - \ell(M/IM) + \ell \left(\widetilde{IM} / IM \right) + \sum_{j=2}^s (j-1)h_j - \ell(H_{\mathfrak{m}}^0(M)), \\ &= e_0^I(M) - \ell(M/IM) + \ell \left(\widetilde{IM} / IM \right) + \sum_{j=2}^s (j-1)h_j - \ell(H_{\mathfrak{m}}^0(M)). \end{aligned}$$

Hence equality holds if and only if $\sum_{j=2}^s (j-1)h_j = 0$ and $\widetilde{IM} = IM$. Assume that equality holds that is, $\widetilde{IM} = IM$ and $\sum_{j=2}^s (j-1)h_j = 0$. We have $h_j = 0$ for $j \geq 2$. So, by graded Nakayama's Lemma one can see that $\widetilde{G}_I(M)$ is generated in degree 0 and 1. This follows that $\widetilde{I^j M} = I^j M + \widetilde{I^{j+1} M}$ for all $j \geq 2$. By Proposition 7.5 we thus get $\widetilde{I^j M} = I^j M + H_{\mathfrak{m}}^0(M)$ for $j \geq 2$. Conversely suppose that $\widetilde{IM} = IM$ and $\widetilde{I^j M} = I^j M + H_{\mathfrak{m}}^0(M)$ for $j \geq 2$. This gives $\widetilde{I^j M} = I^j M + \widetilde{I^{j+1} M}$ for all $j \geq 2$. Which implies that $\widetilde{G}_I(M)$ is generated in degree zero and one. So we have $h_j = 0$ for all $j \geq 2$. $\square$

We give an example which shows that the bound in Theorem 10.6 can be attained.

Example 10.7. Let $R = k[[x, y]]/(xy, y^2)$ with the maximal ideal $\mathfrak{m} = \langle x, y \rangle$. The Hilbert function of R with respect to $\mathfrak{m}$ is

$$\frac{1+z-z^2}{1-z}.$$

We get $e_1(R) - e_0(R) + \ell(R/\mathfrak{m}) = -1$. We now claim $\ell(H_{\mathfrak{m}}^0(R)) = 1$. Let $q = \langle y \rangle$ be the ideal in R . Notice that $\mathfrak{m} \cdot q = 0$. Also $q \neq 0$. Thus $q \simeq k$ as R -modules. Further $R/q \simeq k[[x]]$ is Cohen-Macaulay. Now using the short exact sequence

$$0 \rightarrow q \rightarrow R \rightarrow R/q \rightarrow 0,$$

we get $H_{\mathfrak{m}}^0(R) \simeq H_{\mathfrak{m}}^0(q) = k$.

In view of Theorem 10.6 we pose a question

Question: If $\dim M \geq 2$ then is the set

$$\{ e_1^I(M) - e_0^I(M) + \ell(M/IM) : I \text{ an } \mathfrak{m}\text{-primary ideal} \}$$

bounded below ?

This result holds if M is a generalized Cohen-Macaulay R -module. (see [8, 2.4 and 3.1]).

ACKNOWLEDGMENTS

It is a pleasure to thank Prof. J. K. Verma and Prof A. V. Jayanthan for many discussions. We also thank the referee for many pertinent comments.

REFERENCES

- [1] G. Niesi A. Capani and L. Robbiano, *CoCoA, a system for doing Computations in Commutative Algebra*, 1995, available via anonymous ftp from cocoa.dima.unige.it.
- [2] M. Brodmann, *Asymptotic stability of $\text{Ass}(M/I^n M)$* , Proc. Amer. Math. Soc. **74** (1979), no. 1, 16–18. MR MR521865
- [3] W. Bruns and J. Herzog, *Cohen-Macaulay rings*, Cambridge Studies in Advanced Mathematics, vol. 39, Cambridge University Press, Cambridge, 1993. MR MR1251956
- [4] M. D’Anna, A. Guerrieri, and W. Heinzer, *Invariants of ideals having principal reductions*, Comm. Algebra **29** (2001), 889–906. MR MR1842007 (2002e:13008)
- [5] D. Eisenbud, *Commutative algebra with a view towards algebraic geometry*, Cambridge Studies in Advanced Mathematics, Springer-Verlag, New York, 1991.
- [6] J. Elias, *On the computation of the Ratliff-Rush closure*, J. Symbolic Comput. **37** (2004), no. 6, 717–725. MR MR2095368 (2005j:13022)
- [7] G. Pfister G. M. Gruel and H. Schönemann, *Singular 2.0, a computer algebra systems for polynomial computations*, 2001, Center of Computer Algebra, University of Kaiserslautern, <http://www.singular.uni-kl.de>.
- [8] S. Goto and K. Nishida, *Hilbert coefficients and Buchsbaumness of associated graded rings*, J. Pure Appl. Algebra **181** (2003), 61–74. MR MR1971805
- [9] W. Heinzer, B. Johnston, D. Lantz, and K. Shah, *The Ratliff-Rush ideals in a Noetherian ring: a survey*, Methods in module theory (Colorado Springs, CO, 1991), Lecture Notes in Pure and Appl. Math., vol. 140, Dekker, New York, 1993, pp. 149–159. MR MR1203805
- [10] L. J. Ratliff Jr., *On asymptotic prime divisors*, Pacific J. Math. **111** (1984), no. 2, 395–413. MR MR734863 (85c:13003)
- [11] L. J. Ratliff Jr. and D. E. Rush, *Two notes on reductions of ideals*, Indiana Univ. Math. J. **27** (1978), no. 6, 929–934. MR MR0506202
- [12] S. McAdam, *Asymptotic prime divisors*, Lecture Notes in Mathematics, 1023, Springer-Verlag Berlin Heidelberg, New York Tokyo, 1983. MR MR0485852
- [13] L. Melkersson and P. Schenzel, *Asymptotic prime ideals related to derived functors*, Proc. Amer. Math. Soc. **117** (1993), 935–938. MR MR1124148
- [14] M. Nagata, *Local rings*, Interscience Tracts in Pure and Applied Mathematics, No. 13, Interscience Publishers a division of John Wiley & Sons New York-London, 1962. MR MR0155856
- [15] R. Naghipour, *Ratliff-Rush closure of ideals with respect to a Noetherian module*, Journal of Pure and Applied Algebra **195** (2005), 167–172.
- [16] T. J. Puthenpurakal, *Ratliff-Rush Filtration, Regularity and depth of Higher Associated graded modules: Part I*, Journal of Pure and Applied Algebra **208** (2007), 159–176.
- [17] M. E. Rossi and I. Swanson, *Notes on the behavior of the Ratliff-Rush filtration*, Commutative algebra (Grenoble/Lyon, 2001), Contemp. Math., vol. 331, Amer. Math. Soc., Providence, RI, 2003, pp. 313–328. MR MR2013172
- [18] M. E. Rossi and G. Valla, *Cohen-Macaulay local rings of dimension 2 and an extended version of a conjecture of J. Sally*, J. Pure Appl. Algebra **122** (1997), 293–311.
- [19] ———, *The Hilbert function of the Ratliff-Rush filtration*, J. Pure Appl. Algebra **201** (2005), 25–41. MR MR2158745 (2006g:13008)

- [20] J. D. Sally, *Numbers of generators of ideals in local rings*, Marcel Dekker Inc., New York, 1978. MR MR0485852
- [21] P. Schenzel, *On the use of local cohomology in algebra and geometry*, Six lectures on commutative algebra (Bellaterra, 1996), Progr. Math., vol. 166, Birkhäuser, Basel, 1998, pp. 241–292. MR MR1648667 (99k:13025)

DEPARTMENT OF MATHEMATICS, IIT BOMBAY, POWAI, MUMBAI 400 076, INDIA
E-mail address: `tputhen@math.iitb.ac.in`

DEPARTMENT OF MATHEMATICS, IIT BOMBAY, POWAI, MUMBAI 400 076, INDIA
E-mail address: `fahed@math.iitb.ac.in`